

TWO-DIMENSIONAL QUANTUM POINCARÉ GROUP^{*†}

JAKUB REMBIELIŃSKI[‡]

*Department of Mathematical Physics, University of Łódź
ul. Pomorska 149/153, 90-236 Łódź, Poland*

and

WACŁAW TYBOR[§]

*Department of Theoretical Physics, University of Łódź
ul. Pomorska 149/153, 90-236 Łódź, Poland*

Abstract

Quantum Poincaré-Weyl group in two dimensional quantum Minkowski space-time is considered and an appropriate relativistic kinematics is investigated. It is claimed that a consistent approach to the above questions demands a kind of a “quantum geometry” in the q -deformed space-time.

1 Introduction

Recently, in a number of papers [1, 2, 3, 4, 5, 6, 7, 8] it was considered a variety of models realising the ideas of quantization of space-time and phase-space by means of the non-commutative geometry and quantum group theory methods. In particular in [1] it was considered simple one-dimensional non-commutative dynamical systems. A very intriguing feature of that models is non-commutativity of the inertial mass. However, the non-commutativity of the coordinates with inertial mass is no so satisfactory property because of generically different physical meaning of the coordinates (geometrical objects) and the inertial mass (dynamical object).

In this paper we try to solve simultaneously two problems: (a) the indicated question of the non-commutativity of the inertial mass, (b) the rescaling of the q -Lorentz invariants related to the fact that quantum determinant does not belong to the center of the q -Poincaré group [2]. This can be done by introducing in the q -deformed Minkowski space-time a kind of the “quantum geometry” by means of the notion of “quantum metrics”. In the classical limit the standard Lobatshevsky geometry is recovered. Moreover, in a contrast to the [2] we give a geometric interpretation of momenta as the translation generators defined via corresponding quantum Cartan-Maurer forms.

^{*}Supported by KBN grant No. 2 0218 91 01

[†]Talk given by J.R. during the “Conference on Quantum Topology”, Manhattan, Kansas, 24–28 March 1993.

[‡]e-mail: jaremb@plunlo51.bitnet

[§]e-mail: watybor@plunlo51.bitnet

2 Quantum Minkowski Space-Time and the Quantum Poincaré-Weyl Group

We assume that the q -deformed Minkowski space-time is a *real form* of the Manin's plane. Consequently the q -Minkowski space-time is generated by generators $x_{\pm}$ satisfying

$$x_+ x_- = q x_- x_+, \quad (1)$$

with $x_{\pm}^* = x_{\pm}$ and $|q| = 1$. We will interpret $x_{\pm}$ as light-cone coordinates.

The resulting Poincaré-Weyl group is a subgroup of the $IGL(2)_{q,s,\mu}$ quantum group introduced in [9] with the co-module action of the form

$$\begin{pmatrix} x'_+ \\ x'_- \\ 1 \end{pmatrix} = \delta \begin{pmatrix} x_+ \\ x_- \\ 1 \end{pmatrix} = \begin{pmatrix} \sigma\omega & 0 & u_+ \\ 0 & \sigma\omega^{-1} & u_- \\ 0 & 0 & 1 \end{pmatrix} \otimes \begin{pmatrix} x_+ \\ x_- \\ 1 \end{pmatrix}, \quad (2)$$

where $\sigma^2 = \det_q \begin{pmatrix} \sigma\omega & 0 & u_+ \\ 0 & \sigma\omega^{-1} & u_- \\ 0 & 0 & 1 \end{pmatrix}$ and $\sigma^* = \sigma$, $\omega^* = \omega$, $u_{\pm}^* = u_{\pm}$.

The generators of the above q -Poincaré-Weyl group fulfil the following algebraic rules

$$\omega\sigma = \sigma\omega, \quad (3)$$

$$u_{\pm}\sigma = q^{\pm\frac{1}{2}}\sigma u_{\pm}, \quad (4)$$

$$\omega u_{\pm} = q^{\frac{1}{2}}u_{\pm}\omega, \quad (5)$$

$$u_+ u_- = q u_- u_+. \quad (6)$$

We will follow with the standard Woronowicz form of co-product [10]. The antipode and co-unity have the form

$$g^{-1} = \begin{pmatrix} \sigma^{-1}\omega^{-1} & 0 & -\sigma^{-1}\omega^{-1}u_+ \\ 0 & \sigma^{-1}\omega & -\sigma^{-1}\omega u_- \\ 0 & 0 & 1 \end{pmatrix}; \quad (7)$$

$$\epsilon(\sigma) = \epsilon(\omega) = 1, \quad \epsilon(u_{\pm}) = 0. \quad (8)$$

3 Bicovariant Differential Calculus on Quantum Poincaré-Weyl Group

We can formulate a bicovariant differential calculus on the above quantum group [11] with help of the usual differentials $d\sigma$, $d\omega$, $du_{\pm}$

$$\sigma d\sigma = d\sigma \sigma, \quad (9)$$

$$\sigma d\omega = d\omega \sigma, \quad (10)$$

$$\omega d\omega = d\omega \omega, \quad (11)$$

$$\omega d\sigma = d\sigma \omega, \quad (12)$$

$$u_{\pm} d\omega = q^{-\frac{1}{2}}d\omega u_{\pm}, \quad (13)$$

$$u_{\pm} d\sigma = q^{\pm\frac{1}{2}}d\sigma u_{\pm}, \quad (14)$$

$$\omega du_{\pm} = q^{\frac{1}{2}}du_{\pm} \omega, \quad (15)$$

$$\sigma du_{\pm} = q^{\mp\frac{1}{2}}du_{\pm} \sigma, \quad (16)$$

$$u_{\pm} du_{\pm} = du_{\pm} u_{\pm}, \quad (17)$$

$$u_{\pm} du_{\mp} = q^{\pm 1}du_{\mp} u_{\pm}, \quad (18)$$

$$du_{\pm} d\omega = -q^{-\frac{1}{2}} d\omega du_{\pm}, \quad (19)$$

$$du_{\pm} d\sigma = -q^{\pm\frac{1}{2}} d\sigma du_{\pm}, \quad (20)$$

$$du_+ du_- = -q du_- du_+, \quad (21)$$

$$d\omega d\sigma = -d\sigma d\omega, \quad (22)$$

$$(d\sigma)^2 = (d\omega)^2 = (du_{\pm})^2 = 0. \quad (23)$$

Now we introduce the differential forms

$$\Sigma = \sigma^{-1} d\sigma, \quad \Omega = \omega^{-1} d\omega, \quad T_{\pm} = \sigma^{-1} \omega^{\mp 1} du_{\pm}, \quad (24)$$

with the following commutation relations

$$\Sigma \Omega = -\Omega \Sigma, \quad (25)$$

$$\Sigma T_{\pm} = -T_{\pm} \Sigma, \quad (26)$$

$$\Omega T_{\pm} = -T_{\pm} \Omega, \quad (27)$$

$$T_+ T_- = -q^{-1} T_- T_+. \quad (28)$$

Then we can define, via the Cartan-Maurer form

$$g^{-1} dg \equiv i(D\Sigma + K\Omega + P_+ T_+ + P_- T_-) \quad (29)$$

the generators of the scaling D —dilatation, the Lorentz boost K and translations $P_{\pm}$. Consequently

$$[D, K] = 0, \quad (30)$$

$$[D, P_{\pm}] = -iP_{\pm}, \quad (31)$$

$$[K, P_{\pm}] = \mp iP_{\pm}, \quad (32)$$

$$[P_+, P_-]_{q^{-1}} = 0. \quad (33)$$

Here $[P_+, P_-]_{q^{-1}} = P_+ P_- - q^{-1} P_- P_+$.

4 Covariant Differential Calculus on the q -Minkowski Space-Time and the q -Kinematics

To construct a kind of relativistic kinematics it is necessary to have a covariant differential calculus on the q -Minkowski space. Taking into account the classification of differential calculi for the Manin's plane [12], we obtain

$$x_{\pm} dx_{\pm} = dx_{\pm} x_{\pm}, \quad (34)$$

$$x_{\pm} dx_{\mp} = q^{\pm 1} dx_{\mp} x_{\pm}, \quad (35)$$

$$dx_+ dx_- = -q dx_- dx_+, \quad (36)$$

$$(dx_{\pm})^2 = 0. \quad (37)$$

Now we assume that $x_{\pm}$ depend on an affine parameter τ , i.e.

$$x_{\pm} = x_{\pm}(\tau), \quad dx_{\pm} = \dot{x}_{\pm} d\tau. \quad (38)$$

Consequently, under the existence of the free motion (we admit solutions $\ddot{x}_{\pm} = 0$), we obtain

$$x_{\pm} \dot{x}_{\mp} = q^{\pm 1} \dot{x}_{\mp} x_{\pm}, \quad (39)$$

$$x_{\pm} \dot{x}_{\pm} = \dot{x}_{\pm} x_{\pm}, \quad (40)$$

$$\dot{x}_+ \dot{x}_- = q \dot{x}_- \dot{x}_+. \quad (41)$$

Momentum is defined in a standard way. We assume that the inertial mass m commute with $x_{\pm}$ (i.e. m has not any geometrical meaning).

$$m^* = m, \quad \dot{m} = 0, \quad (42)$$

$$mx_{\pm} = x_{\pm}m. \quad (43)$$

Now for a free particle

$$p_{\pm} = m\dot{x}_{\pm}, \quad (44)$$

so

$$p_{\pm}^* = p_{\pm}, \quad (45)$$

$$p_+p_- = qp_-p_+, \quad (46)$$

according to the geometrical interpretation of $p_{\pm}$.

5 Quantum Geometry

The difference between powers of q in the Eq. (33) and Eq. (46) is related to the contravariant nature of $p_{\pm} = m\dot{x}_{\pm}$ and covariant nature of $P_{\pm}$. This suggests a possibility of an existence of an analogon of the metrics in the q -Minkowski space-time.

Let us define a “qauntum” geometry in our Minkowski space-time by means of a “quantum metrics”; namely

$$ds^2 = (dx_+, \quad dx_-) \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} dx_+ \\ dx_- \end{pmatrix} \quad (47)$$

is replaced by

$$ds_q^2 = (dx_+, \quad dx_-) \begin{pmatrix} 0 & q^{\frac{1}{2}}\Gamma \\ q^{-\frac{1}{2}}\Gamma & 0 \end{pmatrix} \begin{pmatrix} dx_+ \\ dx_- \end{pmatrix}, \quad (48)$$

where the new generator Γ satisfy

$$\Gamma^* = \Gamma, \quad \dot{\Gamma} = 0, \quad \delta(\Gamma) = \sigma^{-2} \otimes \Gamma, \quad (49)$$

$$\Gamma m = m\Gamma, \quad (50)$$

$$\Gamma x_{\pm} = q^{\pm 1} x_{\pm} \Gamma. \quad (51)$$

The “line” element ds_q^2 can be written in a compact form as

$$ds_q^2 = dx^{\mu} g_{\mu\nu} dx^{\nu} \quad (52)$$

with $[g_{\mu\nu}^q] = \begin{pmatrix} 0 & q^{\frac{1}{2}}\Gamma \\ q^{-\frac{1}{2}}\Gamma & 0 \end{pmatrix}$, $\mu, \nu = \pm$. Notice that $[g_q^{\mu\nu}] = \begin{pmatrix} 0 & q^{\frac{1}{2}}\Gamma^{-1} \\ q^{-\frac{1}{2}}\Gamma^{-1} & 0 \end{pmatrix}$.

It is easy to see that ds_q^2 belongs to the center of the space-time algebra. Consequently the square of the relativistic momentum reads

$$p^2 = p^{\mu} g_{\mu\nu} p^{\nu} = q^{\frac{1}{2}} p_+ \Gamma p_- + q^{-\frac{1}{2}} p_- \Gamma p_+ = m^2 C^2, \quad (53)$$

where $C^2 = q^{\frac{1}{2}} \dot{x}_+ \Gamma \dot{x}_- q^{-\frac{1}{2}} + \dot{x}_- \Gamma \dot{x}_+$ is the square of the “light velocity”. Both p^2 and C^2 also belong to the center of the space-time algebra. Note that the relation of momenta satisfying Eq. (33) and Eq. (46) is of the form $P_{\mu} = g_{\mu\nu} p^{\nu}$.

Let us summarize the basic space-time algebra rules

$$x_+x_- = qx_-x_+, \quad (54)$$

$$p_+p_- = qp_-p_+, \quad (55)$$

$$x_{\pm}p_{\pm} = p_{\pm}x_{\pm}, \quad (56)$$

$$x_{\pm}p_{\mp} = q^{\pm 1}p_{\mp}x_{\pm}, \quad (57)$$

$$\Gamma x_{\pm} = q^{\pm 1}x_{\pm}\Gamma, \quad (58)$$

$$\Gamma p_{\pm} = q^{\pm 1}p_{\pm}\Gamma. \quad (59)$$

It is evident that non-commutativity of coordinates and momenta implies that there is a number of “classical” Heisenberg-like uncertainty relations. However, an appropriate analysis demanded a knowledge of representations of the above algebra in the (rigged) Hilbert space framework.

Acknowledgements

We are grateful to K. A. Smoliński for fruitful discussions. One of us (J.R.) thanks to David Yetter for his kind hospitality.

References

- [1] I. Ya. Aref’eva and I. V. Volovich, *Phys. Lett.* **B268** (1991) 179.
- [2] J. Rembieliński, *Phys. Lett.* **B287** (1992) 145.
- [3] J. Rembieliński, in *From the Rotation Group to Quantum Algebras*, ed. B. Gruber (Plenum, New York 1993).
- [4] J. Rembieliński and K. A. Smoliński, Łódź Univ. preprint KFT UŁ 3/93.
- [5] T. Brzeziński, J. Rembieliński and K. A. Smoliński, *Mod. Phys. Lett.* **A8** (1993) 409.
- [6] M. R. Ubriaco, *Mod. Phys. Lett.* **A8** (1993) 89.
- [7] J. Schwenk and J. Wess, *Phys. Lett.* **B291** (1992) 273.
- [8] P. Caban, A. Dobrosielski, A. Krajewska, Z. Walczak, Łódź Univ. preprint KFT UŁ 1/93.
- [9] J. Rembieliński, *Phys. Lett.* **B296** (1992) 335.
- [10] S. L. Woronowicz, *Commun. Math. Phys.* **111** (1987) 613.
- [11] S. L. Woronowicz, *Commun. Math. Phys.* **122** (1989) 125.
- [12] T. Brzeziński, H. Dąbrowski and J. Rembieliński, *J. Math. Phys.* **33** (1992) 19.
- [13] M. Schlieker, M. Weich and R. Weixler, *Z. Phys.* **C53** (1992) 79.